



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification<sup>6</sup> :A61K 38/02, 38/04, 38/16, C07K 7/06,  
7/08

A1

(11) International Publication Number:

WO 98/11907

(43) International Publication Date:

26 March 1998 (26.03.98)

(21) International Application Number: PCT/US97/16217

(22) International Filing Date: 15 September 1997 (15.09.97)

(30) Priority Data:

60/026,978 20 September 1996 (20.09.96) US

08/928,958 12 September 1997 (12.09.97) US

(71) Applicant: BRISTOL-MYERS SQUIBB COMPANY  
[US/US]; Lawrenceville-Princeton Road, Princeton, NJ  
08543-4000 (US).(72) Inventors: NADLER, Steven, G.; 500 W. Roy Street #206W,  
Seattle, WA 98119-3874 (US). CLEAVELAND, Jeffrey,  
S.; 8300 32nd Avenue N.W., Seattle, WA 98107-3921 (US).  
BLAKE, James; 201 Galer Street #243, Seattle, WA 98109-  
3192 (US). HAFFAR, Omar, K.; 101 Elliott Avenue West,  
Seattle, WA 98119 (US).(74) Agents: ROBINS, Roberta, L.; Robins & Associates, Suite 200,  
90 Middlefield Road, Menlo Park, CA 94025 (US) et al.(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR,  
BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE,  
GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO,  
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH,  
KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ,  
BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,  
CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL,  
PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,  
ML, MR, NE, SN, TD, TG).**Published***With international search report.**Before the expiration of the time limit for amending the  
claims and to be republished in the event of the receipt of  
amendments.*(54) Title: PEPTIDE INHIBITORS OF NUCLEAR PROTEIN TRANSLOCATION HAVING NUCLEAR LOCALIZATION SE-  
QUENCES AND METHODS OF USE THEREOF

## (57) Abstract

Novel polypeptide inhibitors of cytoplasmic protein nuclear translocation are disclosed. The inhibitors are comprised of a signal sequence and at least two nuclear localization sequences. The polypeptides are useful as immunosuppression, antiviral and antitumor agents.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

<b>AL</b>	Albania	<b>ES</b>	Spain	<b>LS</b>	Lesotho	<b>SI</b>	Slovenia
<b>AM</b>	Armenia	<b>FI</b>	Finland	<b>LT</b>	Lithuania	<b>SK</b>	Slovakia
<b>AT</b>	Austria	<b>FR</b>	France	<b>LU</b>	Luxembourg	<b>SN</b>	Senegal
<b>AU</b>	Australia	<b>GA</b>	Gabon	<b>LV</b>	Latvia	<b>SZ</b>	Swaziland
<b>AZ</b>	Azerbaijan	<b>GB</b>	United Kingdom	<b>MC</b>	Monaco	<b>TD</b>	Chad
<b>BA</b>	Bosnia and Herzegovina	<b>GE</b>	Georgia	<b>MD</b>	Republic of Moldova	<b>TG</b>	Togo
<b>BB</b>	Barbados	<b>GH</b>	Ghana	<b>MG</b>	Madagascar	<b>TJ</b>	Tajikistan
<b>BE</b>	Belgium	<b>GN</b>	Guinea	<b>MK</b>	The former Yugoslav	<b>TM</b>	Turkmenistan
<b>BF</b>	Burkina Faso	<b>GR</b>	Greece		Republic of Macedonia	<b>TR</b>	Turkey
<b>BG</b>	Bulgaria	<b>HU</b>	Hungary	<b>ML</b>	Mali	<b>TT</b>	Trinidad and Tobago
<b>BJ</b>	Benin	<b>IE</b>	Ireland	<b>MN</b>	Mongolia	<b>UA</b>	Ukraine
<b>BR</b>	Brazil	<b>IL</b>	Israel	<b>MR</b>	Mauritania	<b>UG</b>	Uganda
<b>BY</b>	Belarus	<b>IS</b>	Iceland	<b>MW</b>	Malawi	<b>US</b>	United States of America
<b>CA</b>	Canada	<b>IT</b>	Italy	<b>MX</b>	Mexico	<b>UZ</b>	Uzbekistan
<b>CF</b>	Central African Republic	<b>JP</b>	Japan	<b>NE</b>	Niger	<b>VN</b>	Viet Nam
<b>CG</b>	Congo	<b>KE</b>	Kenya	<b>NL</b>	Netherlands	<b>YU</b>	Yugoslavia
<b>CH</b>	Switzerland	<b>KG</b>	Kyrgyzstan	<b>NO</b>	Norway	<b>ZW</b>	Zimbabwe
<b>CI</b>	Côte d'Ivoire	<b>KP</b>	Democratic People's	<b>NZ</b>	New Zealand		
<b>CM</b>	Cameroon		Republic of Korea	<b>PL</b>	Poland		
<b>CN</b>	China	<b>KR</b>	Republic of Korea	<b>PT</b>	Portugal		
<b>CU</b>	Cuba	<b>KZ</b>	Kazakhstan	<b>RO</b>	Romania		
<b>CZ</b>	Czech Republic	<b>LC</b>	Saint Lucia	<b>RU</b>	Russian Federation		
<b>DE</b>	Germany	<b>LI</b>	Liechtenstein	<b>SD</b>	Sudan		
<b>DK</b>	Denmark	<b>LK</b>	Sri Lanka	<b>SE</b>	Sweden		
<b>EE</b>	Estonia	<b>LR</b>	Liberia	<b>SG</b>	Singapore		

5     PEPTIDE INHIBITORS OF NUCLEAR PROTEIN TRANSLOCATION  
          HAVING NUCLEAR LOCALIZATION SEQUENCES  
          AND METHODS OF USE THEREOF

Technical Field

10           The present invention relates generally to  
polypeptide inhibitors of gene expression. More  
particularly, the invention relates to polypeptide  
inhibitors of nuclear protein translocation which have  
gene expression modulating activity, immunosuppressive  
15 activity, antiviral activity, and antitumor activity.

Background of the Invention

          Nuclear transport is essential to a number  
of biological processes including gene expression and  
20 cell division, as well as to viral replication,  
tumorigenesis and tumor cell proliferation. The  
mechanism of nuclear transport has only recently been  
characterized in detail and has been shown to involve  
a number of discrete steps. Proteins that are  
25 destined to be transported into the nucleus contain  
within their amino acid sequence a short stretch of  
amino acids termed a nuclear localization sequence  
("NLS"). These sequences are generally basic in  
nature, however, there has been no consensus sequence  
30 identified. Thus, there is a wide variety of these  
sequences that appear to be specific for particular  
proteins.

          Within the cell, these NLSs may be either  
masked or unmasked by accessory proteins or by  
35 conformational changes within the NLS-containing  
protein. An NLS may be masked because it is buried in  
the core of the protein and not exposed on the surface

of the protein. Unmasking of NLSs, and nuclear translocation of cytoplasmic proteins may be triggered by phosphorylation, dephosphorylation, proteolytic digestion, subunit association or dissociation of an inhibitory subunit, or the like. Accordingly, the masking and unmasking of NLSs provides a mechanism by which the transport of these cytoplasmic proteins into the nucleus may be regulated.

Nuclear translocation of transcription factors requires the presence of an unmasked or activated NLS in the nucleus-targeted protein. The binding of certain ligands to cell surface receptors activates the nuclear translocation of cytoplasmic transcription factors. Once in the nucleus, these transcription factors exert gene expression modulatory activity.

NF- $\kappa$ B is a ubiquitous transcription factor found in various levels and states of activation in different cell types. NF- $\kappa$ B is composed of several different subunits including p65, p50, c-rel, p52 and p105. Recent studies suggest that distinct NF- $\kappa$ B complexes contribute to the regulatory control of gene transcription. The function and regulation of NF- $\kappa$ B has been most well-characterized in lymphocytic cells. In these cells, there is a wide variety of target genes, e.g., immunoregulatory genes, that are regulated by NF- $\kappa$ B including  $\kappa$  Ig light chains. Such genes include those that encode the interleukin-2 $\alpha$  ("IL-2 $\alpha$ ") receptor, interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), tumor necrosis factor- $\alpha$  ("TNF- $\alpha$ "), and the like.

In unstimulated cells, a major form of NF- $\kappa$ B is a heterodimer of p50 and p65 (RelA) subunits. Nonactive NF- $\kappa$ B is retained in the cytoplasm as an inactive complex by inhibitory proteins such as I $\kappa$ B $\alpha$ ,  $\beta$  and  $\gamma$ . When cells are appropriately stimulated, e.g., by a proinflammatory stimulus such as a

cytokine, the I $\kappa$ Bs are degraded, thereby releasing free NF- $\kappa$ B dimers, which translocate to the nucleus and activate target genes, e.g., lymphokine genes and other immunoregulatory genes. This response is  
5 transient and is terminated through delayed NF- $\kappa$ B-mediated I $\kappa$ B $\alpha$  induction.

Recently it has been demonstrated that glucocorticoids exert their immunosuppressive activity by inhibiting NF- $\kappa$ B nuclear translocation. Scheinman  
10 et al. (1995) *Science* 270:283-286 and Auphan et al. (1995) *Science* 270:286-290 independently demonstrated that the inhibition is mediated by an increase in the induction by glucocorticoids of I $\kappa$ B inhibitory proteins. These investigators proposed that  
15 inhibitors of NF- $\kappa$ B may be useful immunosuppressive and anti-inflammatory agents. Such an NF- $\kappa$ B nuclear translocation inhibitor, comprising an NLS from the p50 subunit of NF- $\kappa$ B attached to a membrane-permeable polypeptide motif, was described in Lin et al. (1995)  
20 *J. Biol. Chem.* 270:14255-14258.

Nuclear translocation of proteins other than endogenous transcription factors and other cytoplasmic proteins also depends on the presence of an activated or unmasked NLS. For example, nuclear translocation  
25 of the retroviral preintegration complex is a crucial step in human immunodeficiency virus type-1 ("HIV-1") replication in nondividing cells such as monocytes and growth-arrested T cells. Such translocation is dependent on the presence of an NLS in the N-terminal  
30 portion of HIV matrix antigen ("MA") p17. Indeed, the HIV-1 enhancer contains tandem binding sites for NF- $\kappa$ B that can be essential for virus replication (Ross et al. (1991) *J. Virol.* 65:4350-4358; Parrott et al. (1991) *J. Virol.* 65:1414-1419). Nuclear translocation  
35 of the HIV-1 preintegration complex can be partially inhibited by an excess of the SV40 large T antigen NLS (Gulizia et al. (1994) *J. Virol.* 68:2021-2025).

Furthermore, Dubrovsky et al. (1995) *Molecular Med.* 2:217-230 reported that a series of compounds capable of binding to and reacting with the HIV-1 MA p17 NLS inhibit HIV-1 replication in human monocytes.

5           In addition, tumorigenesis and tumor cell proliferation are regulated by the expression of oncoproteins, many of which are cytoplasmic transcription factors that are translocated into the nucleus by virtue of the presence of an NLS. Miller  
10 et al. (1996) *J. Cell Biochemistry* 60:560.

          Accordingly, inhibitors of nuclear translocation of cytoplasmic proteins would be useful as gene expression modulating agents, immunoregulatory agents, antiviral agents, antitumor agents, and the  
15 like.

#### Summary of the Invention

          The present invention provides for a polypeptide that can be introduced into an intact cell  
20 for the purpose of inhibiting the nuclear translocation of a cytoplasmic protein. The polypeptide contains at least two NLSs and an amino acid sequence that can deliver the polypeptide through the cytoplasmic membrane into the cell. The inventors  
25 herein have found that such a polypeptide exhibits surprisingly superior characteristics compared to a polypeptide having only one NLS.

          Accordingly, in one embodiment, the invention is directed to a polypeptide comprising a  
30 signal sequence peptide and at least two NLSs covalently attached thereto.

          In another embodiment, the invention is directed to a method of introducing an exogenous polypeptide comprising an NLS into an intact cell to  
35 inhibit nuclear translocation of a cellular protein. The method includes providing a polypeptide comprising a signal sequence peptide and at least two NLSs as

described above, and contacting the cell with the polypeptide for a period of time effective to introduce the exogenous polypeptide into the cell.

5 In still another embodiment, the invention is directed to a method of suppressing an immune response of a subject comprising administering to the subject an immunosuppressive amount of a polypeptide comprising a signal sequence peptide and at least two NLSs.

10 In a further embodiment, the invention is directed to a method of treating or preventing a viral infection in an individual comprising administering to the individual an effective antiviral amount of a polypeptide inhibitor of nuclear translocation of a  
15 cellular protein, said inhibitor comprising a signal sequence peptide and at least two NLSs.

In yet a further embodiment, the invention is directed to a method of transcriptionally modulating the expression of cellular genes comprising  
20 contacting a target cell with an inhibitor of nuclear translocation of a cellular protein, wherein the inhibitor comprises a signal sequence peptide and at least two NLSs.

25 These and other embodiments of the subject invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

#### Brief Description of the Figures

30 FIG. 1A is a graphical representation of the effect of PKKKRKVA VALLPAVLLALLAPKKRKVC (SEQ ID NO:1) (the "SV40MEM" polypeptide) on lipopolysaccharide ("LPS")-stimulated surface antigen expression in 70Z/3 murine leukemia pre-B cells (solid bar:  $\kappa$  Ig light chain; stippled bar: IL-2 $\alpha$  receptor; striped bar:  
35 CD45). FIG. 1B is a graphical representation of the effect of three inhibitory polypeptides on LPS-stimulated  $\kappa$  Ig light chain production in 70Z/3 murine

leukemia pre-B cells (diamonds: the SV40MEM polypeptide; squares: KKKYKAAVALLPAVLLALLAKKKYKC (SEQ ID NO:2) (the "HIV-1MEM" polypeptide); triangles: AKRVKLA VALLPAVLLALLAKRVKLC (SEQ ID NO:3) (the "C-MYCMEM" polypeptide)). FIG. 1C is a graphical representation of the effect of increasing LPS concentrations on SV40MEM-inhibited  $\kappa$  Ig light chain production in 70Z/3 murine leukemia pre-B cells.

FIG. 2 is a graphical representation of the effect of the SV40MEM polypeptide on LPS-stimulated cytokine production in 70Z/3 murine leukemia pre-B cells (squares: untreated; diamonds: the SV40MEM polypeptide (10  $\mu$ M); circles: the SV40MEM polypeptide (5  $\mu$ M); triangles: the SV40 NLS (10  $\mu$ M)).

FIG. 3 is a graphical representation of dose-response relationships of three inhibitory polypeptides on LPS-stimulated  $\kappa$  Ig light chain production in 70Z/3 murine leukemia pre-B cells (triangles: a peptide containing only the fibroblast growth factor ("FGF") signal sequence AAVALLPAVLLALLAP (SEQ ID NO:4) (the "MEM" peptide); squares: a polypeptide having the FGF signal sequence and the NLS of NF- $\kappa$ B p50 on the carboxy terminus of the signal sequence, namely, AAVALLPAVLLALLAPVQRKRQKLMP (SEQ ID NO:5) polypeptide (the "NF- $\kappa$ B MEM" polypeptide); diamonds: the SV40MEM polypeptide comprised of L-amino acids; circles: the SV40MEM polypeptide comprised of D-amino acids.).

FIG. 4A is a graphical representation of the effect of inhibitory polypeptides on LPS-stimulated TNF- $\alpha$  production. FIG. 4B is a graphical representation of the effect of inhibitory polypeptides on LPS-stimulated interleukin-8 ("IL-8") production. In both FIG. 4A and FIG. 4B the following symbols are used: squares - media control; crossed squares - the MEM peptide (5  $\mu$ M); diamonds - the NF- $\kappa$ B MEM polypeptide (5  $\mu$ M); circles - the SV40MEM



polypeptide comprised of L-amino acids (5  $\mu$ M); and triangles - the SV40MEM polypeptide comprised of D-amino acids (5  $\mu$ M).

5 **FIG. 5** is a graphical representation of the effect of the SV40MEM polypeptide on LPS-induced CD40 expression in 70Z/3 murine leukemia pre-B cells.

**FIG. 6A** is a graphical representation of the effect of the HIV-1MEM polypeptide on  $^3$ H-deoxyribothymidine uptake into peripheral blood  
10 mononuclear cells ("PBMCs"). **FIG. 6B** is a graphical representation of the effect of HIV-1MEM on viral p24 production in anti-CD3 stimulated PBMCs infected with HIV-1 primary isolate M1. **FIG. 6C** is a photograph of a gel depicting the effect of HIV-1MEM polypeptide on  
15 the expression of proviral gag sequences in anti-CD3 activated PBMCs infected with HIV-1 primary isolate M1.

**FIG. 7** is a photograph of a gel depicting the results of a polymerase chain reaction analysis of  
20 the effect of HIV-1MEM polypeptide on the expression of proviral gag sequences in H9 human lymphoma T-cells or Jurkat human leukemia T-cells infected with HIV-1 primary isolate M1.

**FIG. 8A** is a photograph of gels depicting the results of polymerase chain analyses of the effect  
25 of HIV-1MEM and the NF- $\kappa$ BMEM polypeptide on the expression of proviral gag sequences in Jurkat T-cells infected with HIV<sub>LAI</sub>. **FIG. 8B** is a photograph of gels depicting the results of polymerase chain analyses of  
30 the effect of HIV-1MEM and the NF- $\kappa$ BMEM polypeptide on the expression of 2-long terminal repeat ("LTR") circles in Jurkat T-cells infected with HIV<sub>LAI</sub>.

**FIG. 9** is a graphical representation of the effect of SV40MEM prepared from D-amino acids on  
35 proliferation of 70Z/3 murine leukemia pre-B cells.

FIG. 10 is a graphical representation of the effect of SV40MEM prepared from D-amino acids on proliferation of RAJI human B-cell leukemia cell line.

FIG. 11A is a graphical representation of the effect of an intravenous administration of the SV40MEM polypeptide on the *in vivo* response of mice to sheep red blood cells. FIG. 11B is a graphical representation of the effect of an oral administration of the SV40MEM polypeptide on the *in vivo* response of mice to sheep red blood cells.

FIG. 12 shows the effect of BMS 205820 and C-MYCMEM (BMS 214572) on the anti-hemocyanin (KLH) response in mice.

FIG. 13A, shows the effect of the BMS 205820 polypeptide on the production of TNF- $\alpha$  *in vivo*. FIG. 13B shows the effect of the BMS 205820 polypeptide on the production of IL-6 *in vivo*. FIG. 13C shows the effect of the BMS 205820 polypeptide on the production of IL-10 *in vivo*.

FIG. 14 depicts the effect of BMS 205820, C-MYCMEM (BMS 214572), an SV40 NLS polypeptide containing a single NLS, a c-myc NLS alone, without a translocation sequence (AKRVKL (SEQ ID NO:6)) and a control non-NLS polypeptide, 377G, on lipopolysaccharide (LPS) binding to CD14.

FIG. 15A is a time line depicting the administration scheme for the intraperitoneal injection of ovalbumin (OVA), nebulized OVA, BMS 205820 and C-MYCMEM. FIG. 15B shows the % eosinophils in lung following the treatment outlined in FIG. 15A.

#### Detailed Description

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of protein chemistry and biochemistry, molecular biology, microbiology and recombinant DNA technology, which are within the skill of the art.

Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989); *DNA Cloning*, Vols. I and II (D.N. Glover ed. 1985);  
5 *Oligonucleotide Synthesis* (M.J. Gait ed. 1984);  
*Nucleic Acid Hybridization* (B.D. Hames & S.J. Higgins eds. 1984); *Animal Cell Culture* (R.K. Freshney ed. 1986); *Immobilized Cells and Enzymes* (IRL press, 1986); Perbal, B., *A Practical Guide to Molecular*  
10 *Cloning* (1984); the series, *Methods In Enzymology* (S. Colowick and N. Kaplan eds., Academic Press, Inc.).

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural references unless the content  
15 clearly dictates otherwise.

#### A. Definitions

In describing the present invention, the following terms will be employed, and are intended to  
20 be defined as indicated below.

As used herein, the term "signal sequence" or "signal sequence peptide" is used to indicate a peptide that is capable of directing the movement of the polypeptide of which it is a part through a cell  
25 membrane. In particular, the term is used to indicate a peptide that directs the movement of a polypeptide across the cytoplasmic membrane into the cell. The term "signal sequence" is intended to encompass not only the signal sequence of a particular polypeptide,  
30 but also fragments or derivatives thereof that are capable of delivering a polypeptide through a cell membrane. A "signal sequence" may be composed of L- or D-amino acids.

The terms "nuclear localization sequence" and "NLS" are used interchangeably to indicate a  
35 peptide that directs the transport of a protein with which it is associated from the cytoplasm of a cell

across the nuclear envelope barrier. The term "NLS" is intended to encompass not only the nuclear localization sequence of a particular peptide, but also derivatives thereof that are capable of directing translocation of a cytoplasmic polypeptide across the nuclear envelope barrier. NLSs are capable of directing nuclear translocation of a polypeptide when attached to the N-terminus, the C-terminus, or both the N- and C- termini of the polypeptide. In addition, a polypeptide having an NLS coupled by its N- or C-terminus to amino acid side chains located randomly along the amino acid sequence of the polypeptide will be translocated. Adam et al. (1990) *J. Cell. Biol.* 111:807-818. "Nuclear localization sequences" may be composed of D- or L-amino acids.

By "interchangeably flanked at its amino- and carboxy-termini by a first and a second NLS" is intended to mean that the first or second NLS may be located at either the amino- or carboxy-terminus of the signal sequence polypeptide.

An "inhibitor of nuclear translocation" is a polypeptide composed of a signal sequence peptide and at least two NLSs which inhibits, e.g., either decreases or halts, nuclear localization of a cytoplasmic protein. Preferably, the polypeptide comprises a signal sequence peptide interchangeably flanked at its amino- and carboxy-termini by a first and a second NLS. The NLSs at the N- and C-termini may be the same or different. In one preferred embodiment, the signal sequence peptide and the NLSs are each composed of L-amino acids. In another preferred embodiment, the signal sequence peptide and the NLSs are each composed of D-amino acids.

A "derivative" of a polypeptide is intended to include homologous polypeptides in which conservative amino acid substitutions have been made, as well as to include other amino acid substitutions

that result in a polypeptide that retains its function, e.g., as a signal sequence peptide, an NLS, or an inhibitor of nuclear localization. A "derivative" of a peptide may be a peptide mimetic.

5           "Peptide mimetics" are structures which serve as substitutes for peptides in interactions with acceptor molecules (see Morgan et al. (1989) *Ann. Reports Med. Chem.* 24:243-252 for a review of peptide mimetics). Peptide mimetics, as used herein, include  
10       synthetic structures which may or may not contain amino acids and/or peptide bonds, but retain the structural and functional features of a peptide ligand. The term, "peptide mimetics" also includes  
15       peptoids and oligopeptoids, which are peptides or oligomers of N-substituted amino acids (Simon et al. (1972) *Proc. Natl. Acad. Sci. USA* 89:9367-9371). Further included as peptide mimetics are peptide  
20       libraries, which are collections of peptides designed to be of a given amino acid length and representing all conceivable sequences of amino acids corresponding thereto. Methods for the production of peptide  
      mimetics are described more fully below.

      Two polypeptide sequences are "substantially homologous" when at least about 85% (preferably at  
25       least about 85% to 90%, and most preferably at least about 95%) of the nucleotides or amino acids match over a defined length of the molecule. As used herein, substantially homologous also refers to sequences showing identity to the specified  
30       polypeptide sequence.

      The terms "polypeptide", "peptide" and "protein" are used interchangeably and refer to any polymer of amino acids (dipeptide or greater) linked through peptide bonds. Thus, the terms "polypeptide",  
35       "peptide" and "protein" include oligopeptides, protein fragments, analogues, muteins, fusion proteins and the like.

The following single-letter amino acid abbreviations are used throughout the text:

	Alanine	A	Arginine	R
	Asparagine	N	Aspartic acid	D
5	Cysteine	C	Glutamine	Q
	Glutamic acid	E	Glycine	G
	Histidine	H	Isoleucine	I
	Leucine	L	Lysine	K
	Methionine	M	Phenylalanine	F
10	Proline	P	Serine	S
	Threonine	T	Tryptophan	W
	Tyrosine	Y	Valine	V

By an "isolated polypeptide" is meant a polypeptide which is devoid of, in whole or part, tissue or cellular components with which the protein is normally associated in nature. Thus, a polypeptide contained in a tissue extract would constitute an "isolated" polypeptide, as would a polypeptide synthetically or recombinantly produced.

By "mammalian subject" is meant any member of the class Mammalia, including, without limitation, humans and non-human primates, such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; and laboratory animals including rodents such as mice, rats and guinea pigs. The term does not denote a particular age. Thus, adult, newborn and fetal mammals are intended to be covered.

The term "treatment" as used herein refers to either (i) the prevention of infection or reinfection (prophylaxis), or (ii) the reduction or elimination of symptoms of the disease of interest (therapy).

## B. General Methods

Central to the present invention is the discovery of polypeptide molecules that inhibit nuclear localization of cytoplasmic proteins. These molecules comprise a signal sequence peptide and at least two NLSs. The polypeptide inhibitors, and derivatives thereof, provide useful tools for introducing an exogenous polypeptide comprising an NLS into an intact cell to inhibit nuclear translocation of a cellular protein, for studying the role of nuclear translocation in the regulation of cellular processes.

In addition, since the nuclear translocation of certain cellular peptides is required for the host organism to mount an immune response, the polypeptide inhibitors are useful as immunosuppression agents. Immune responses are typically manifested by the expression of antibodies, the production of a number of cytokines, and/or the expression of cell surface receptors. Thus, inhibition of immune responses by the inhibitory peptides can take the form of: inhibition of antibody production, including the production of antibody component peptides such as a  $\kappa$  light chain polypeptide; inhibition of cytokine production, including such cytokines as interleukin-1, interleukin-2, interleukin-4, interleukin-6, interleukin-10, tumor necrosis factor, or granulocyte-macrophage colony-stimulating factor; and/or the inhibition of the expression of cell-surface receptors such as an interleukin-2 receptor, gp39, CD40, CD45, CD80, CD86, ICAM, ELAM, major histocompatibility complex ("MHC") class II, or VCAM. Clark et al. (1994) *Nature* 367:425.

By virtue of their immunosuppressive activities, the polypeptide inhibitors of the present invention are useful in the treatment of a wide variety of immune disorders, including but not limited

to, the treatment of autoimmune diseases such as rheumatoid arthritis, multiple sclerosis, juvenile-onset diabetes, systemic lupus erythematosus (SLE), autoimmune uveoretinitis, autoimmune vasculitis, 5 bullous pemphigus, myasthenia gravis, autoimmune thyroiditis or Hashimoto's disease, Sjogren's syndrome, granulomatous orchitis, autoimmune oophoritis, Crohn's disease, sarcoidosis, rheumatic carditis, ankylosing spondylitis, Grave's disease, and 10 autoimmune thrombocytopenic purpura (see e.g., Paul, W.E. (1993) *Fundamental Immunology*, Third Edition, Raven Press, New York, Chapter 30, pp. 1033-1097; and Cohen et al. (1994) *Autoimmune Disease Models, A Guidebook*, Academic Press, 1994).

15 Similarly, the polypeptide inhibitors of the present invention are useful for treating physical symptoms manifested by responses to allergens which can initiate a state of hypersensitivity, or which can provoke a hypersensitivity reaction in a subject 20 already sensitized with the allergen. Such physical symptoms include asthma, joint swelling, urticaria, and the like. Additionally, due to their immunosuppressive properties, the polypeptide inhibitors of the present invention are useful in the 25 treatment of sepsis and in the prevention of septic shock, a potentially lethal condition caused by the uncontrolled production of certain cytokines due to the presence of endotoxins, such as lipopolysaccharide (LPS), from extracellular bacteria.

30 Furthermore, since many viruses, e.g., herpes virus, cytomegalovirus, retroviruses, and the like, make use of the host cell's nuclear translocation machinery, the inhibitory polypeptides are useful as antiviral agents. In addition, since 35 tumorigenesis and tumor cell proliferation appear to be mediated by the expression of oncogenes to make oncoproteins, many of which are transcription factors



that are translocated into the nucleus, Miller et al. (1996), *supra*, the polypeptide inhibitors, or derivatives thereof, can be used to suppress tumor growth.

5           The claimed inhibitors include sequences of amino acids that comprise signal sequences from such polypeptides as the antennapedia homeodomain, FGF, HIV Tat, or Hsc70, and derivatives or mimetics thereof capable of delivering the inhibitor through the  
10   cytoplasmic membrane into the cell. Preferred signal sequences include RQIKIWFQNRRMKWKK (SEQ ID NO:7), AAVALLPAVLLALLA (SEQ ID NO:8), AAVALLPAVLLALLAP (SEQ ID NO:4), CFITKALGISYGRKKRRQRRRPPQGSQTH (SEQ ID NO:9), and the like, or derivatives or mimetics thereof  
15   capable of delivering the inhibitor through the cytoplasmic membrane into the cell.

          Candidate signal sequences can be tested for their ability to direct the translocation of proteins across cell membranes, for example, by monitoring the  
20   localization of exogenous detectably labeled proteins into the cell cytoplasm. Lin et al. (1995), *supra*, describe the use of radiolabeled proteins. *In vitro* nuclear peptide import can be measured using NLS peptides coupled to a fluorescent protein by methods  
25   described in Adam et al. (1990), *supra*.

          The polypeptide inhibitor further comprises at least two NLSs. The NLSs can be covalently bonded to the N-terminus, to the C-terminus, to both the N- and C- termini of the signal sequence polypeptide, to  
30   amino acid side chains located along the amino acid sequence of the signal sequence polypeptide, or any combination thereof. Preferably, the signal sequence polypeptide is interchangeably flanked at its amino- and carboxy-termini by a first and a second NLS. The  
35   first and second NLSs may be the same or different. A discussion of NLSs and a list of NLSs can be found in

Boulikas (1993) *Crit. Rev. Eukaryotic Gene Expression* 3:193-227, and references cited therein.

Approaches for identifying NLSs include: (1) gene fusion experiments between a candidate NLS-coding DNA segment and the gene coding for a cytoplasmic protein (see, e.g., Silver et al. (1984) *Proc. Natl. Acad. Sci. U.S.A.* 81:5951; Moreland et al. (1987) *Mol. Cell. Biol.* 7:4048; and Picard et al. (1987) *EMBO J.* 6:3333); (2) nuclear import of nonnuclear proteins conjugated to synthetic NLS peptides (see, e.g., Goldfarb et al. (1986) *Nature* 322:641; Markland et al. (1987) *Mol. Cell. Biol.* 7:4255; and Chelsky et al. (1989) *Mol. Cell. Biol.* 9:2487); and (3) site-directed mutagenesis of a specific segment of a nuclear protein, resulting in its cytoplasmic retention (see, e.g., Greenspan et al. (1988) *J. Virol.* 62:3020; van Etten et al. (1989) *Cell* 58:669; and Boulukos et al. (1989) *Mol. Cell. Biol.* 9:5718).

Preferred NLSs include PKKKRKV (SEQ ID NO:10) and KKKRKVC (SEQ ID NO:11) from the SV40 large T antigen (see, Kalderon et al. (1984) *Cell* 39:499), GKRSKA (SEQ ID NO:12) from yeast histone H2B (see, Moreland et al. (1987) *Mol. Cell. Biol.* 7:4048), KRPRP (SEQ ID NO:13) from adenovirus E1A (see, Lyons et al. (1987) *Mol. Cell. Biol.* 7:2451), GNKAKRQRST (SEQ ID NO:14) from the v-rel oncogene of the avian reticuloendotheliosis retrovirus strain T (see, Gilmore et al. (1988) *J. Virol.* 62:703), GGAAKRVKLD (SEQ ID NO:15) from the human c-myc oncoprotein (see, Chelsky et al. (1989) *Mol. Cell. Biol.* 9:2487), SALIKKKKKMAP (SEQ ID NO:16) from the murine c-abl (IV) gene product (see, Van Etten et al. (1989) *Cell* 58:669), RKLKKLGN (SEQ ID NO:17) from the human or rat androgen receptor (see, Guiochon-Mantel et al. (1989) *Cell* 57:1147), PQPKKKP (SEQ ID NO:18) from protein p53 (see, Dang et al. (1989) *J. Biol. Chem.* 264:18019), ASKSRKRKL (SEQ ID NO:19) from viral Jun, a

transcription factor of the AP-1 complex (see, Chida et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:4290), KKKYK (SEQ ID NO:20) and KKKYKC (SEQ ID NO:21), both of which are from the human immunodeficiency virus matrix protein (see, Bukrinsky et al. (1993) *Nature* 365:666), KSKKK (SEQ ID NO:22) from the human immunodeficiency virus matrix 2 protein (see, Bukrinsky et al. (1993), *supra*), AKRVKL (SEQ ID NO:6) and KRVKLC (SEQ ID NO:23) both of which are from the human *c-myc* oncoprotein (see, Chelsky et al. (1989), *supra*), and derivatives and mimetics thereof that are effective as an NLS.

The polypeptide inhibitors of the present invention may be synthesized by conventional techniques known in the art, for example, by chemical synthesis such as solid phase peptide synthesis. Such methods are known to those skilled in the art. In general, these methods employ either solid or solution phase synthesis methods, well known in the art. See, e.g., J. M. Stewart and J. D. Young, *Solid Phase Peptide Synthesis*, 2nd Ed., Pierce Chemical Co., Rockford, IL (1984) and G. Barany and R. B. Merrifield, *The Peptides: Analysis, Synthesis, Biology*, editors E. Gross and J. Meienhofer, Vol. 2, Academic Press, New York, (1980), pp. 3-254, for solid phase peptide synthesis techniques; and M. Bodansky, *Principles of Peptide Synthesis*, Springer-Verlag, Berlin (1984) and E. Gross and J. Meienhofer, Eds., *The Peptides: Analysis, Synthesis, Biology*, *supra*, Vol. 1, for classical solution synthesis.

In general, these methods comprise the sequential addition of one or more amino acids or suitably protected amino acids to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid is protected by a suitable protecting group. The protected or derivatized amino acid can then be either attached to an inert solid

support or utilized in solution by adding the next amino acid in the sequence having the complementary (amino or carboxyl) group suitably protected, under conditions suitable for forming the amide linkage.

5 The protecting group is then removed from this newly added amino acid residue and the next amino acid (suitably protected) is then added, and so forth. After all the desired amino acids have been linked in the proper sequence, any remaining protecting groups  
10 and any solid support are removed either sequentially or concurrently to afford the final polypeptide. By simple modification of this general procedure, it is possible to add more than one amino acid at a time to a growing chain, for example, by coupling (under  
15 condition that do not racemize chiral centers) a protected tripeptide with a properly protected dipeptide to form, after deprotection, a pentapeptide.

Typical protecting groups include t-butyloxycarbonyl (Boc), 9-fluorenylmethoxycarbonyl  
20 (Fmoc), benzyloxycarbonyl (Cbz), p-toluenesulfonyl (Tos); 2,4-dinitrophenyl, benzyl (Bzl), biphenyl-isopropyloxycarboxycarbonyl, cyclohexyl, isopropyl, acetyl, o-nitrophenylsulfonyl, and the like. Of these, Boc and Fmoc are preferred.

25 Typical solid supports are generally cross-linked polymeric materials. These include divinylbenzene cross-linked styrene-based polymers, for example, divinylbenzene-hydroxymethylstyrene copolymers, divinylbenzene-chloromethylstyrene  
30 copolymers, and divinylbenzene-benzhydrylaminopolystyrene copolymers. The Divinylbenzene-benzhydrylaminopolystyrene copolymers, as illustrated herein using p-methyl-benzhydrylamine resin, offers the advantage of directly introducing a  
35 terminal amide functional group into the peptide chain, which function is retained by the chain when the chain is cleaved from the support.

In one preferred method, the polypeptides are prepared by conventional solid phase chemical synthesis on, for example, an Applied Biosystems, Inc. (ABI) 430A peptide synthesizer using a resin that  
5 permits the synthesis of the amide peptide form and using t-Boc amino acid derivatives (Peninsula Laboratories, Inc.) with standard solvents and reagents. Polypeptides containing either L- or D-amino acids may be synthesized in this manner. Polypeptide  
10 composition is confirmed by quantitative amino acid analysis and the specific sequence of each peptide may be determined by sequence analysis.

Alternatively, the polypeptides can be produced by recombinant DNA techniques by synthesizing  
15 DNA encoding the desired polypeptide, along with an ATG initiation codon. Once coding sequences for the desired polypeptides have been synthesized or isolated, they can be cloned into any suitable vector for expression. Numerous cloning vectors are known to  
20 those of skill in the art, and the selection of an appropriate cloning vector is a matter of choice. Examples of recombinant DNA vectors for cloning and host cells which they can transform include the bacteriophage  $\lambda$  (*E. coli*), pBR322 (*E. coli*), pACYC177  
25 (*E. coli*), pKT230 (gram-negative bacteria), pGV1106 (gram-negative bacteria), pLAFR1 (gram-negative bacteria), pME290 (non-*E. coli* gram-negative bacteria), pHV14 (*E. coli* and *Bacillus subtilis*), pBD9 (*Bacillus*), pIJ61 (*Streptomyces*), pUC6 (*Streptomyces*),  
30 YIp5 (*Saccharomyces*), YCp19 (*Saccharomyces*) and bovine papilloma virus (mammalian cells). See, generally, *DNA Cloning: Vols. I & II, supra*; Sambrook et al., *supra*; B. Perbal, *supra*. Insect cell expression systems, such as baculovirus systems, can also be used  
35 and are known to those of skill in the art and described in, e.g., Summers and Smith, *Texas Agricultural Experiment Station Bulletin No. 1555*

(1987). Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *inter alia*, Invitrogen, San Diego CA ("MaxBac" kit).

5           The gene can be placed under the control of a promoter, ribosome binding site (for bacterial expression) and, optionally, an operator (collectively referred to herein as "control" elements), so that the DNA sequence encoding the desired polypeptide is  
10 transcribed into RNA in the host cell transformed by a vector containing this expression construction. The coding sequence may or may not contain a signal peptide or leader sequence. Heterologous leader sequences can be added to the coding sequence which  
15 cause the secretion of the expressed polypeptide from the host organism. Leader sequences can be removed by the host in post-translational processing. See, e.g., U.S. Patent Nos. 4,431,739; 4,425,437; 4,338,397.

          Other regulatory sequences may also be  
20 desirable which allow for regulation of expression of the protein sequences relative to the growth of the host cell. Such regulatory sequences are known to those of skill in the art, and examples include those which cause the expression of a gene to be turned on  
25 or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Other types of regulatory elements may also be present in the vector, for example, enhancer sequences.

          The control sequences and other regulatory  
30 sequences may be ligated to the coding sequence prior to insertion into a vector, such as the cloning vectors described above. Alternatively, the coding sequence can be cloned directly into an expression vector which already contains the control sequences  
35 and an appropriate restriction site.

          In some cases it may be necessary to modify the coding sequence so that it may be attached to the

control sequences with the appropriate orientation; i.e., to maintain the proper reading frame. It may also be desirable to produce mutants or analogs of the polypeptide of interest. Mutants or analogs may be prepared by the deletion of a portion of the sequence encoding the protein, by insertion of a sequence, and/or by substitution of one or more nucleotides within the sequence. Techniques for modifying nucleotide sequences, such as site-directed mutagenesis, are well known to those skilled in the art. See, e.g., Sambrook et al., *supra*; *DNA Cloning*, Vols. I and II, *supra*; *Nucleic Acid Hybridization*, *supra*.

The expression vector is then used to transform an appropriate host cell. A number of mammalian cell lines are known in the art and include immortalized cell lines available from the American Type Culture Collection (ATCC), such as, but not limited to, Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), Madin-Darby bovine kidney ("MDBK") cells, as well as others. Similarly, bacterial hosts such as *E. coli*, *Bacillus subtilis*, and *Streptococcus* spp., will find use with the present expression constructs. Yeast hosts useful in the present invention include *inter alia*, *Saccharomyces cerevisiae*, *Candida albicans*, *Candida maltosa*, *Hansenula polymorpha*, *Kluyveromyces fragilis*, *Kluyveromyces lactis*, *Pichia guillerimondii*, *Pichia pastoris*, *Schizosaccharomyces pombe* and *Yarrowia lipolytica*. Insect cells for use with baculovirus expression vectors include, *inter alia*, *Aedes aegypti*, *Autographa californica*, *Bombyx mori*, *Drosophila melanogaster*, *Spodoptera frugiperda*, and *Trichoplusia ni*. The proteins may also be expressed in *Trypanosomes*.

Depending on the expression system and host selected, the proteins of the present invention are produced by growing host cells transformed by an expression vector described above under conditions whereby the protein of interest is expressed. The protein is then isolated from the host cells and purified. If the expression system secretes the protein into growth media, the protein can be purified directly from the media. If the protein is not secreted, it is isolated from cell lysates. The selection of the appropriate growth conditions and recovery methods are within the skill of the art. Once purified, the amino acid sequences of the proteins can be determined, i.e., by repetitive cycles of Edman degradation, followed by amino acid analysis by HPLC. Other methods of amino acid sequencing are also known in the art.

As explained above, peptide mimetics which structurally and functionally mimic the peptide inhibitors described above will also find use herein and may be generated using the following strategies and procedures. Generally, mimetics are designed based on information obtained by systematic replacement of L-amino acids by D-amino acids or, in the case of a polypeptide inhibitor that is made of D-amino acids, the systematic replacement of D-amino acids by L-amino acids, replacement of side chain moieties by a methyl group or pseudoisosteric groups with different electronic properties (see Hruby et al. (1990) *Biochem. J.* 268:249-262), and by systematic replacement of peptide bonds in the above described polypeptide inhibitors with amide bond replacements. For example, analogues containing amide bond surrogates may be used to investigate aspects of peptide structure and function, such as rotational freedom in the backbone, intra- and intermolecular hydrogen-bond patterns, modifications of local and



total polarity and hydrophobicity, and oral bioavailability.

Local conformational constraints can also be introduced to determine conformational requirements for activity of a candidate polypeptide mimetic inhibitor of nuclear translocation. For example,  $\beta,\beta$ -disubstituted amino acids may be used to examine the effects of conformational constraints on peptide activity (see, e.g., Manning et al. (1982) *J. Med. Chem.* 25:408-414; Mosberg et al. (1983) *Proc. Natl. Acad. Sci. USA* 106:506-512; Pelton et al. (1985) *Proc. Natl. Acad. Sci. USA* 82:236-239).

The mimetics can include isosteric amide bonds such as  $\Psi[\text{CH}_2\text{S}]$ ,  $\Psi[\text{CH}_2\text{NH}]$ ,  $\Psi[\text{CSNH}_2]$ ,  $\Psi[\text{NHCO}]$ ,  $\Psi[\text{COCH}_2]$  and  $\Psi[(\text{E}) \text{ or } (\text{Z}) \text{ CH}=\text{CH}]$  (see, for review, Spatola (1983) in "Chemistry and Biochemistry of Amino Acids, Peptides and Proteins," Volume VII, (Weinstein, ed.), Marcel Dekker, New York, 267-357). Structures which mimic the tetrahedral transition state associated with hydrolysis of a substrate bond can also be present and include hydroxymethylene, fluoroketone moieties and phosphoramidate transition state mimics (Bühlmayer et al. (1988) *J. Med. Chem.* 31:1839; Sham et al. (1988) *FEBS Lett.* 220:299; Matthews (1988) *Acc. Chem. Res.* 21:333). Cyclic amino acid analogues may be used to constrain amino acid residues to particular conformational states, e.g.,  $\alpha\alpha'$ - and  $\beta\beta'$ -substituted cyclic amino acids such as 1-aminocyclopentanecarboxylic acid (cycloleucine) and  $\beta,\beta$ -cyclopentamethylene- $\beta$ -mercaptopropionic acid (see Hruby et al. (1990), *supra*).

The mimetics can also include mimics of inhibitor peptide secondary structure -- structures which can model the 3-dimensional orientation of amino acid residues into the known secondary conformations of proteins -- including  $\beta$ -turn mimetics, such as phenoxathin ring system, and  $\beta$ -sheet mimics, such as

epindolidione structures. Design, synthesis and conformational analysis of an  $\alpha$ -helix inducing template has been described (Kemp et al. (1988) *Tetrahedron Lett.* 29:4931; Kemp et al. (1988) *Tetrahedron Lett.* 29:4935). Similarly, peptoids will find use herein. Peptoids are oligomers of N-substituted amino acids (Simon et al. (1972), *supra*), and can be used as motifs for the generation of chemically diverse libraries of novel molecules, which can then be tested for nuclear translocation inhibitory activity. The monomers may incorporate *t*-butyl-based side-chain and 9-fluorenylmethoxy-carbonyl  $\alpha$ -amine protection. Oligomerization of the peptoid monomers may be performed by, for example, *in situ* activation by either benzotriazol-1-yloxytris(pyrrolidino)phosphonium hexafluorophosphate or bromotris(pyrrolidino)phosphonium hexafluorophosphate. Other steps are identical to conventional peptide synthesis using  $\alpha$ -(9-fluorenylmethoxycarbonyl)amino acids. Oligopeptoids may be identified which have activities comparable to the corresponding inhibitory polypeptides and, thus, are useful as inhibitors of nuclear translocation (see Simon et al. (1992), *supra*).

Peptide ligands that exhibit nuclear translocation inhibitory activity can be developed by using a biological expression system (see Christian et al. (1992) *J. Mol. Biol.* 227:711-8; Devlin et al. (1990) *Science* 249:404-406; Cwirla et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6378-6382). The use of such systems allows the production of large libraries of random peptide sequences and the screening of these libraries for peptide sequences that have desired biochemical activity. The libraries may be produced by cloning synthetic DNA that encodes random peptide sequences into *Escherichia coli* expression vectors. In the filamentous phage system, foreign peptide

sequences can be expressed on the surface of the infectious phage (see Smith (1985) *Science* 228:1315-1317; Parmley et al. (1988) *Gene* 73:305-318).

For example, a library may be made by  
5 ligating into an appropriate phage, a synthetic DNA fragment containing a degenerate coding sequence  $(NNK)_n$ , where N stands for an equal mixture of the deoxynucleotides G, A, T, and C, K stands for an equimolar mixture of G and T, and n stands for the  
10 number of amino acid residues desired in the product peptide. Phage are screened for expression of inhibitory activity. Those that express inhibitory activity can be cloned and propagated, their DNAs sequenced to determine the amino acid sequences of  
15 their expressed polypeptide which can be assessed for their ability to inhibit nuclear translocation.

Large libraries of polypeptide inhibitors can also be constructed by concurrent synthesis of overlapping peptides as described in U.S. Patent No.  
20 4,708,871 to Geysen. The solid support is generally a polyethylene or polypropylene rod onto which is graft polymerized a vinyl monomer containing at least one functional group to produce polymeric chains on the carrier. The functional groups are reacted to provide  
25 primary or secondary amine groups which are sequentially reacted with amino acid residues in the appropriate order to build the desired synthetic peptide using conventional methods of solid phase peptide chemistry.

30 Once synthesized or otherwise produced, the inhibitory activity of a candidate polypeptide or polypeptide mimetic can be tested by assessing the ability of the candidate to inhibit the lipopolysaccharide-induced nuclear translocation of  
35 NF- $\kappa$ B by, for example, using murine endothelial cells by the method described in Lin et al. (1995), *supra*.

Inhibitory polypeptides can be used in pharmaceutical compositions to treat autoimmune diseases and viral infections, or to suppress transplant rejection. In addition, the polypeptides  
5 can be administered to cancerous tissues in order to suppress tumor growth. The inhibitory polypeptides of the present invention can be formulated into therapeutic compositions in a variety of dosage forms such as, but not limited to, liquid solutions or  
10 suspensions, tablets, pills, powders, suppositories, polymeric microcapsules or microvesicles, liposomes, and injectable or infusible solutions. The preferred form depends upon the mode of administration and the particular cancer type targeted. The compositions  
15 also preferably include pharmaceutically acceptable vehicles, carriers or adjuvants, well known in the art, such as human serum albumin, ion exchangers, alumina, lecithin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate,  
20 and salts or electrolytes such as protamine sulfate. Suitable vehicles are, for example, water, saline, dextrose, glycerol, ethanol, or the like, and combinations thereof. Actual methods of preparing such compositions are known, or will be apparent, to  
25 those skilled in the art. See, e.g., *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Easton, Pennsylvania, 18th edition, 1990.

The above compositions can be administered using conventional modes of delivery including, but  
30 not limited to, intravenous, intraperitoneal, oral, intralymphatic, or subcutaneous administration. Local administration to a tumor in question, or to a site of inflammation, e.g., direct injection into an arthritic joint, will also find use with the present invention.

35 Therapeutically effective doses will be easily determined by one of skill in the art and will depend on the severity and course of the disease, the

patient's health and response to treatment, and the judgment of the treating physician.

C. Experimental

5                   Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

10                   Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

15                                   Experimental Methods

Peptide Preparation: In general, polypeptides can be synthesized using a stepwise solid-phase synthesis method and purified on a C<sub>18</sub> reverse phase high  
20 performance liquid chromatography on a reverse-phase column eluted with a linear gradient of 10%-60% acetonitrile, 0.1% trifluoroacetic acid as described in Merrifield (1963) *J. Am. Chem. Soc.* 85:2149-2154 and Lin et al. (1988) *Biochemistry* 27:5640-5645.

25                   The peptides described herein were synthesized on a Gilson AMS-422 multiple peptide synthesizer using Fmoc amino acids. Gausepohl et al. (1992) *Peptide Res.* 5:315-320. Peptides were cleaved from the resin and deprotected by a 2-hr reaction with  
30 trifluoroacetic acid/water/thioanisole/ethanedithiol (100:5:5:2.5). the peptides were precipitated from ether, redissolved in formic acid, diluted with water, and lyophilized. Purification was achieved by reverse-phase high performance liquid chromatography  
35 using a gradient of acetonitrile in 0.1% trifluoroacetic acid. All synthetic peptides were

characterized by mass spectroscopy on a Bio-Ion 20 instrument and gave the expected molecular weight.

Murine 70Z/3 pre-B leukemia cells (ATCC T1D-158), H9 human lymphoma T-cells (ATCC CRL-8543),  
5 Jurkat human leukemia T-cells (ATCC T1B-152), and THP-1 human monocytic cells (ATCC T1B-202) were obtained from the American Type Culture Collection.

Expression of  $\kappa$  Ig, CD45, CD40, and the interleukin-2 receptor was assayed using standard  
10 enzyme-linked immunosorbent assays ("ELISAs") (see, Coligan et al. (eds.) (1991) Current Protocols in Immunology, Wiley & Sons, p. 2.10 or by FACS staining to measure cell-surface markers as described in Raff (1970) *Immunology* 19:637). Antibodies form cell  
15 surface markers were obtained from Pharmingen Co.

Amounts of TNF- $\alpha$ , IL-6, IL-8 and IL-10 were determined using the ELISA kits obtained from Genzyme following the manufacturer's instructions.

HIV-1 M1 p 24 levels were assayed using the  
20 procedures described in Smithgall et al (1995) *AIDS Res. and Human Retrovirus* 11:885. Proviral DNA content (gag) was measured using polymerase chain reaction analysis as described in Smithgall et al. (1995), *supra*.

25 2-LTR circles, a strictly nuclear form of HIV-1 DNA the expression of which indicates nuclear localization of the viral genome, was measured by as described in Bukrinsky et al (1991) *Science* 254:423.

30 Example 1  
Inhibition by the SV40MEM Polypeptide of  
 $\kappa$  Ig Light Chain Expression

This experiment was conducted to show that the polypeptide PKKKRKVAVALLPVLLALLAPKKKRKVC (SEQ ID  
35 NO:1) (the "SV40MEM" polypeptide), which comprises the hydrophobic region of the signal peptide from fibroblast growth factor and the SV40 large T antigen

NLS, has immunosuppressive activity. The effect of the SV40MEM polypeptide was compared with two additional polypeptides, KKKYKAAVALLPAVLLALLAKKKYKC (SEQ ID NO:2) (the "HIV-1MEM" polypeptide), which is  
5 composed of the membrane translocation domain of FGF flanked by the NLS from the HIV-1 matrix protein and AKRVKLAAVALLPAVLLALLAKRVKLC (SEQ ID NO:3) (the "C-MYCMEM" polypeptide), which contains the NLS from the human c-myc oncoprotein (Chelsky et al. (1989), *supra*,  
10 and Dang et al. (1988), *supra*) using murine 70Z/3 pre-B cells.

Murine 70Z/3 cells (ATCC T1D-158) are a model cell line for analyzing the effects of compounds on B-cell differentiation. In response to *S. typhosa*  
15 LPS (Difco) or  $\gamma$ -interferon, these cells differentiate from surface  $\kappa$  immunoglobulin negative to  $\kappa$  Ig positive.

As seen in **FIG. 1A**, the SV40MEM polypeptide caused an approximately 75-80% inhibition of  $\kappa$  light  
20 chain expression in response to *S. typhosa* LPS (Difco) after a 1-hr pretreatment with the SV40MEM polypeptide. There was no effect of SV40MEM on the expression of CD45 or the interleukin-2 receptor. The maximum inhibitory effect occurred with an  
25 approximately 3-hr SV40MEM pretreatment. This inhibitory effect of the SV40MEM polypeptide can be overcome by increasing the concentration of LPS.

**FIG. 1B** depicts dose-response data for the effect of the D-amino acid form of the SV40MEM  
30 polypeptide compared with L-amino acid forms of the HIV-1MEM polypeptide and AKRVKLAAVALLPAVLLALLAKRVKLC (SEQ ID NO:3). These data show that the three polypeptides are approximately equally efficacious but that the latter two polypeptides are not as potent at  
35 inhibiting  $\kappa$  Ig light chain production as is SV40MEM. **FIG. 1C** shows that increasing the concentration of LPS can overcome the inhibitory effect of SV40MEM.

Example 2Inhibition by the SV40MEM Polypeptide of  
Cytokine Production

THP-1 control cells were harvested, washed  
5 and set up in RPMI 1640, 10% fetal calf serum, at a  
concentration of  $2.56 \times 10^6$  cells/ml. Phosphate  
buffered saline ("PBS") or one of the two respective  
polypeptides SV40MEM or SV40 (a peptide having only  
10 the SV40 large T antigen NLS sequence) were added to  
the cells at a concentration of 10  $\mu$ M or 20  $\mu$ M. The  
cells were then plated into a 96-well plate with 0.1  
ml/well. The plate was then incubated for 2 hr at  
37°C in a 5% CO<sub>2</sub> atmosphere. Following the initial  
incubation, media or a titration of LPS was added to  
15 each of the wells in a volume of 0.1 ml. Thus, the  
final concentrations of polypeptides were 5  $\mu$ M or 10  
 $\mu$ M. Following 5 hr of LPS stimulation, supernatants  
were taken and TNF- $\alpha$  and IL-8 levels were determined  
by ELISA. In addition, cells treated with PBS or  
20 polypeptides, but not incubated with LPS, were removed  
at the time of supernatant removal to determine the  
viability of the cells following 7 hr of contact with  
the polypeptides. From the data depicted in FIG. 2,  
it can be seen that the SV40 polypeptide had no effect  
25 on TNF- $\alpha$  production but that the SV40MEM polypeptide  
inhibited TNF- $\alpha$  production in a dose-dependent manner.  
Similar results were obtained with respect to IL-8  
production (data not shown).

30

Example 3Comparison of the Inhibition by the SV40MEM  
Polypeptide and other Inhibitory Polypeptides  
of  $\kappa$  Iq Light Chain Expression

This experiment was conducted to compare the  
35 inhibitory effect of the SV40MEM polypeptide with that  
of the polypeptide AAVALLPAVLLALLAPVQRKRQKLMP (SEQ ID



NO:5) (the "NF- $\kappa$ B MEM" polypeptide) on  $\kappa$  light chain expression in 70Z/3 B-cells.

The 70Z/3 cells were pretreated for 2 hr with between 0 and 50  $\mu$ M of: (1) a peptide containing only the FGF signal sequence AAVALLPAVLLALLAP (SEQ ID NO:4) (the "MEM" peptide); (2) a polypeptide having the FGF signal sequence and the NLS of NF- $\kappa$ B p50 on the carboxy terminus of the signal sequence, namely, the NF- $\kappa$ B MEM polypeptide; (3) the SV40MEM polypeptide comprised of L-amino acids; or (4) the SV40MEM polypeptide comprised of D-amino acids. The cells were then incubated with LPS as described in Example 2 and assayed for expression of  $\kappa$  light chain expression.

The data in FIG. 3 show that the L- and D-amino acid forms of the SV40MEM polypeptide were equally potent and efficacious with respect to inhibiting LPS-stimulated  $\kappa$  Ig expression. In addition, the data show that both the L- and D-amino acid forms of the SV40MEM polypeptide were more potent and more efficacious than the NF- $\kappa$ B MEM polypeptide having a single NLS sequence. The EC<sub>50</sub> for both the L- and D-amino acid forms of the SV40MEM polypeptide was 0.7  $\mu$ M, while that for the NF- $\kappa$ B MEM polypeptide having a single NLS sequence was 32  $\mu$ M.

#### Example 4

##### Comparison of the Inhibition by the SV40MEM Polypeptide Other Inhibitory Polypeptides of Cytokine Production

This experiment was conducted to compare the inhibitory effect of the SV40MEM polypeptide and other inhibitory polypeptides on LPS induction of cytokine production. THP-1 control cells (ATCC 9/95 stock) were harvested, washed and set up in RPMI 1640, 10% fetal calf serum as described in Example B. The cells were incubated for 2 hr with PBS or (1) the FGF signal

sequence, (2) a polypeptide having the FGF signal sequence and the NLS of NF- $\kappa$ B p50 on the carboxy terminus of the signal sequence, (3) the SV40MEM polypeptide comprised of L-amino acids, or (4) the SV40MEM polypeptide comprised of D-amino acids at a final concentration of 5  $\mu$ M. The cells were then plated into a 96-well plate with 0.1 ml/well. The plate was then incubated for 2 hr at 37°C in a 5% CO<sub>2</sub> atmosphere. Following the initial incubation, media or a titration of LPS was added to each of the wells in a volume of 0.01 ml/well. Following 5 hr of LPS stimulation, supernatants were taken and TNF- $\alpha$  levels were determined by ELISA. The data depicted in FIG. 4A indicated that only the L- or D-amino acid forms of the SV40MEM polypeptide significantly inhibited LPS-induced TNF- $\alpha$  expression; i.e., the polypeptide comprising the FGF signal sequence peptide and the NF- $\kappa$ B NLS did not significantly inhibit TNF- $\alpha$  expression. At high LPS concentrations, TNF- $\alpha$  was induced even in cells that had been incubated with the L- or D-amino acid forms of the SV40MEM polypeptide. Similar results were obtained with LPS-induced IL-8 production (FIG. 4B).

25

#### Example 5

##### Effect of the SV40MEM Polypeptide on CD40 Expression

70Z/3 murine B-cells were treated with the SV40MEM polypeptide for 3 hr prior to activation with LPS. After activation with LPS, there was an approximately four-fold increase in CD40 expression above basal levels. The SV40MEM polypeptide inhibited this up-regulation of CD40 (see FIG. 5). These results indicate that the SV40MEM polypeptide should be efficacious at inhibiting B-cell responses in vivo.

35

Example 6Inhibition of Infection of Peripheral Blood  
Mononuclear Cells with the Primary Isolate  
of HIV-1 M1 by the HIV-1MEM Polypeptide

5 PBMCs were obtained washed, cultured, and  
infected with HIV-1 as described in Dubrovsky et al.  
(1995) *Mol. Med.* 1:217-230). PBMCs from a  
seronegative donor were collected and depleted of CD8<sup>+</sup>  
T cells by negative selection. The cells were  
10 activated with an anti-CD3 monoclonal antibody  
prepared according to the method described in Wee et  
al. (1993) *J. Exp. Med.* 177:219 and incubated for 3 hr  
with the HIV-1MEM polypeptide. The cells were then  
incubated with M1 virus isolated from PBMC collected  
15 from HIV-infected donors in the presence of inhibitors  
for 2 hr. The virus was washed out, and the cultures  
supplemented with the appropriate inhibitor and  
allowed to incubate for seven days. Virus production  
was determined by measuring the p24 levels in the  
20 culture supernatant (**FIG. 6B**) and contrasted with the  
effect of the polypeptide on cell proliferation (**FIG.**  
**6A**), which virus load was determined by polymerase  
chain reaction analysis of proviral DNA content (gag)  
(**FIG. 6C**). The MEM peptide represents the FGF  
25 membrane translocating portion of the HIV-1MEM  
polypeptide. The HIV-1MEM polypeptide significantly  
reduced the virus load and virus production compared  
to untreated but activated cultures.

30

Example 7Inhibition of Infection of T Cells  
with HIV<sub>M1</sub> Virus by the HIV-1MEM Polypeptide

H9 and Jurkat cells were incubated with the  
HIV-1MEM polypeptide (2  $\mu$ M) or the control MEM peptide  
35 for 4 hr prior to addition of HIV-M1 virus. Cells  
were harvested on day 3 following injection and the  
level of HIV-1 DNA evaluated by polymerase chain

reaction. As shown in FIG. 7, the HIV-1MEM polypeptide inhibited infection of both cells lines as determined by the lower levels of HIV-1 DNA compared to untreated and MEM peptide-treated cultures.

5

#### Example 8

##### Reduction in Viral Infection Due to Reduction in Nuclear Translocation of HIV Genome

Jurkat cells were incubated with the SV40MEM polypeptide for 3 hr with HIV<sub>LAI</sub> for 1-2 hr. The virus was washed out and the cultures supplemented with the appropriate inhibitor (SV40MEM, 0.1  $\mu$ M and 1.0  $\mu$ M; NF- $\kappa$ BMEM, 10  $\mu$ M). After 24 hr, the cells were collected, lysed and evaluated for HIV-1 DNA by polymerase chain reaction analysis of gag sequences (see FIG. 8A) and 2-LTR circles (see FIG. 8B). The uniform signal achieved in all samples when assayed for gag sequences shows that all cultures internalized virus relatively equally and allowed formation of the viral cDNA in the cytoplasm. The difference in the levels of 2-LTR circles shows that SV40MEM inhibited translocation of the viral genome into the nucleus. Thus, the polypeptides do not affect virus entry into cells but, as predicted, specifically target nuclear translocation. The NF- $\kappa$ BMEM peptide showed no effect on viral internalization or translocation of the viral genome into the nucleus.

#### Example 9

##### Effect of SV40MEM on Proliferation of 70Z/3 Murine B-cell Leukemia Cells

Murine 70Z/3 B-cells were treated with 0, 0.2, 0.5, 1, or 2  $\mu$ M SV40MEM prepared from D-amino acids. On day 0, the cell concentrations of the control and treated cells were the same. After 48 hr of exposure to the peptide, the treated cells were again counted and compared to control cells. FIG. 9

shows that the proliferation of the murine leukemia cell line was inhibited by the lowest concentration (0.2  $\mu$ M) of D-SV40MEM tested and was 90% inhibited by the highest concentration tested (2  $\mu$ M).

5

#### Example 10

##### Effect of SV40MEM on Proliferation of RAJI Human B-cell Leukemia Cells

Human RAJI B-cells were treated with 0, 5,  
10 10 or 25  $\mu$ M SV40MEM prepared from D-amino acids. On  
day 0, the cell concentrations of the control and  
treated cells were the same. After 48 hr of exposure  
to the peptide, the treated cells were again counted  
and compared to control cells. FIG. 10 shows that the  
15 proliferation of the murine leukemia cell line was  
inhibited by the lowest concentration (5  $\mu$ M) of D-  
SV40MEM tested and was greater than 95% inhibited by  
the highest concentration tested (25  $\mu$ M).

20

#### Example 11

##### In Vivo Immunosuppressive Effect of the SV40MEM Polypeptide In a Murine Sheep Red Blood Cell Assay

Female BALB/c mice (6-8 weeks of age) were  
housed in groups of 5 in Thorne units with *ad libitum*  
25 access to food and water. Mice were immunized on day  
0 by intravenous (IV) injection of  $1 \times 10^8$  sheep red  
blood cells ("SRBCs"). The mice were then divided  
into groups of five and control solutions or  
inhibitory peptides were administered at various  
30 doses, either by IV injection or orally ("PO") by  
gavage with a 24-gauge feeding tube and routes.

The SV40MEM polypeptide was administered  
either intravenously (IV) or orally (PO) with the  
following schedule: IV administration of 5 mg/kg on  
35 days 0, 1, 2, and 3; PO administration of 1 mg/kg on  
days 0, 1, 2, 3, and 4. After 14 days, the mice were  
bled from the orbital sinus and the level of anti-SRBC

antibodies were measured in the serum by ELISA. Briefly, Immulon 2 plates (Dynatech) were coated with SRBC membrane antigen (Anal. Biochem. 82:362-371 (1978)) at 5  $\mu$ g/ml in 0.05 M carbonate/bicarbonate buffer (pH 9.6) and incubated overnight at 2-8°C. Plates were blocked for 1-hr at room temperature with Specimen Diluent Concentrate (Genetic Systems, Redmond, WA) diluted 1:10 with distilled water. The plates were washed in phosphate-buffered saline and 0.05% Tween-20 ("PBS/Tween"). Serum samples were diluted in Specimen Diluent and incubated for 1 hr at room temperature. After washing the plates in PBS/Tween, goat anti-mouse IgG1-horse radish peroxidase-conjugated ("IgG1-HRP") antibody (Southern Biotechnology, Birmingham AL) was added at 1:5000 dilution and incubated for 1 hr at room temperature. Plates were washed and conjugate binding was detected using 3,3',5,5'-tetramethylbenzidine ("TMB") (Genetic Systems) as the substrate. TMB was added and incubated for 15 min at room temperature. The reaction was stopped by the addition of 3 N H<sub>2</sub>SO<sub>4</sub>. Absorbance at 45 nm was recorded from duplicate samples using a microtiter plate reader (Bio Tek Instruments).

The results presented in FIG. 11A and FIG. 11B show that, while there was no effect on total IgG levels in the mice receiving the SV40MEM polypeptide either IV or PO, the polypeptide inhibited the specific anti-SRBC response when administered either IV or PO.

#### Example 12

#### Comparison of the Inhibition by the BMS 205820 Polypeptide and other Inhibitory Polypeptides of the Anti-Hemocyanin Response in Mice

Female BALB/c mice (8 weeks of age) were immunized by intraperitoneal injection of 25  $\mu$ g

keyhole limpet (*Megathura crenulata*) hemocyanin ("KLH") (Pacific Bio-Marine Laboratories, Venice, CA) without adjuvant on day 0. Mice were divided into groups of five and inhibitory peptides C-MYCMEM and  
5 BMS 205820 (PKKKRKVAAVALLPAVLLALLAPKKRKV (SEQ ID NO:24)), which is identical to the SV40MEM polypeptide with the exception that the polypeptide lacks the C-terminal cysteine, were administered on day 0 at 5 mg/kg, followed by 5 subsequent doses of the same  
10 concentration every other day. Control mice were treated with a single 200- $\mu$ g dose of Chimeric L6 antibody ("Chi L6"). After 14 days, the mice were bled from the orbital sinus and the level of anti-KLH antibodies were measured in the serum by ELISA as  
15 described in Example 11.

Both the BMS 205820 and the C-MYCMEM polypeptide inhibited the specific anti-KLH response (see FIG. 12).

20

### Example 13

#### Stimulation of Apoptosis by the SV40MEM Polypeptide in Anti-CD3-treated Jurkat T-Cells

A 24-hr incubation of Jurkat T-cells with SV40MEM and anti-CD3, caused greater than 85% of the  
25 cells to apoptose. Neither SV40MEM nor anti-CD3 alone caused significant apoptosis.

### Example 14

#### Inhibition of I $\kappa$ B Degradation by the SV40MEM Polypeptide

30

I $\kappa$ B is an inhibitor of NF- $\kappa$ B nuclear translocation. When cells are activated by the appropriate proinflammatory stimulus, e.g., cytokines or LPS, I $\kappa$ B is degraded. This results in the  
35 unmasking of the NLS of NF- $\kappa$ B, thereby allowing NF- $\kappa$ B to be translocated into the nucleus. This experiment

was done to explore the effect of the SV40MEM polypeptide on I $\kappa$ B degradation in murine 70Z/3 cells.

Murine 70Z/3 cells ( $5 \times 10^7$  cells) were incubated with the SV40MEM polypeptide ( $2 \mu\text{M}$ ) or PBS. The two groups of cells were subdivided into aliquots and activated with LPS ( $100 \text{ ng/ml}$ ) for 0, 15, 30, 60, or 120 min. The cells were lysed in  $200 \mu\text{L}$  lysing buffer ( $20 \text{ mM}$  N-(2-hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid) ("HEPES"), pH 7.2,  $20 \text{ mM}$  NaCl,  $2.5 \text{ mM}$   $\text{MgCl}_2$ ,  $0.1\%$  nonylphenoxy polyethoxy ethanol ("NP-40")). The lysates were frozen until analyzed for the presence of I $\kappa$ B by Western analysis using an antibody directed to the carboxy terminus of I $\kappa$ B (Santa Cruz Corp., Santa Cruz, CA) according to the manufacturer's instructions.

Degradation of I $\kappa$ B was apparent after 15-min activation of the control cells with LPS. There was no apparent degradation of I $\kappa$ B in cells pretreated with the SV40MEM peptide.

#### Example 15

##### Ability of BMS 205820 to Treat Septic Shock

These experiments show that BMS 205820 is efficacious for the treatment of septic shock. A lethal model of septic shock was used as follows.

A. Balb/c mice were pretreated with either  $5 \text{ mg/kg}$  BMS 205820 or PBS, intravenously at 1 and 3 hours, prior to injection with  $200 \text{ mg}$  lipopolysaccharide (LPS), the causative agent of septic shock. LPS was administered intraperitoneally. Mice were given another injection of the polypeptide immediately after the LPS injection, as well as 24 hours after the LPS injection. Mice typically died within 48 hours of the LPS injection. As can be seen in Table 1, in two separate experiments, administration of BMS 205820 significantly increased



the number of mice surviving seven days post- LPS injection.

Table 1

5

Fraction of Mice Surviving Seven  
Days Post LPS Injection

<u>Experiment</u>	<u>LPS + PBS</u>	<u>LPS + BMS 205820</u>
1	0/10	6/10
2	2/10	10/10

10

B. Balb/c were treated with 200  $\mu$ g of LPS followed by the BMS 205820 polypeptide, at the times indicated in Table 2. The BMS polypeptide was administered intravenously and the LPS was

15 administered intraperitoneally. Group A mice were treated with 5 mg/kg of polypeptide in PBS at the indicated times, post- LPS administration. Group B mice received an additional dose of polypeptide 3 hours post-LPS treatment.

20

As can be seen in Table 2, BMS 205820 was effective in inhibiting death in response to LPS, especially when two doses of polypeptide were administered.

25

Table 2

Effect of Delayed Treatment with BMS 205820  
in a Lethal Model of Septic Shock

	<u>Treatment</u>	<u>Fraction Surviving</u>	
		<u>Group A</u>	<u>Group B</u>
30	None	0/5	2/5
	T=0	5/5	5/5
	T=30 min.	1/5	5/5
	T=1 hr.	2/5	5/5
	T=1.5 hr.	1/5	4/5
35	T=2 hr.	0/5	0/5

Example 16Inhibition of Cytokine Production in vivo  
by the BMS 205820 Polypeptide

These experiments show that the BMS 205820  
5 polypeptide is capable of modulating cytokine  
production. In particular, a non-lethal model of  
septic shock was used. Balb/c mice were treated with  
1  $\mu$ g LPS, and at the same time, a single dose of 3  
mg/kg of the BMS 205820 polypeptide in PBS. TNF- $\alpha$ ,  
10 IL-6 and IL-10 levels were measured in serum by ELISA.

As shown in FIG. 13A, the BMS 205820  
polypeptide caused a significant inhibition in the  
production of TNF- $\alpha$ , which is a major contributor to  
the sepsis syndrome. IL-6 levels were only slightly  
15 affected (FIG. 13B), whereas IL-10 production was  
greatly increased (FIG. 13C). The effect on IL-10 is  
beneficial since this cytokine is immunosuppressive  
and has been shown to have efficacy in models of  
septic shock.

20

Example 17Inhibition of LPS Binding to CD14 Using  
BMS 205820 and C-MYCMEM

These experiments show that BMS 205820 and  
25 C-MYCMEM (the "BMS 214572" polypeptide) are both  
capable of inhibiting the binding of LPS to its cell  
surface receptor, CD14, evidencing that the  
polypeptide inhibitors are useful in the treatment of  
sepsis.

30 96-well plates were coated with goat-anti  
human Ig, soluble CD-14 Ig was added, and the plates  
were incubated overnight at 4°C. The BMS polypeptide,  
the C-MYCMEM polypeptide, an SV40 NLS polypeptide  
containing a single SV40 large T-antigen NLS, a c-myc  
35 NLS without a translocation sequence (AKRVKL (SEQ ID  
NO:6)), a control non-NLS polypeptide, 377G, or PBS,  
were added to the plates and the plates were incubated

for 5 min. *E. coli* AO16 LPS was then added to the plates and the plates were incubated for 2 hours at 4°C. The plates were washed and LPS was detected using 0.5 µg/ml mouse anti-*E. coli* LPS, followed by goat anti-mouse horseradish peroxidase.

**FIG. 14** shows the results of the experiment. Both BMS 205820 and C-MYCMEM were able to bind to LPS and block it from binding CD14. The c-myc NLS did not block binding of LPS to CD14.

LPS is the causative agent of septic shock. Thus, by blocking binding of LPS to CD14, the NLS polypeptides described herein may serve to prevent or diminish septic shock *in vivo*. This result was unexpected. Without being bound by a particular theory, it is possible that positively charged NLS polypeptides bind to negatively charged regions on LPS.

#### Example 18

##### Ability of BMS 205820 to Treat Asthma

These experiments show that the BMS 205820 polypeptide is efficacious for the treatment of asthma. The BMS 205820 polypeptide was tested in a mouse model of asthma (Renz et al. (1992) *J. Allergy Clin. Path.* 89:1127-1138). Balb/c mice were administered ovalbumin in an alum adjuvant intraperitoneally at various times, as indicated in **FIG. 15A**. Treatment was followed by nebulization of ovalbumin into the lungs, also as indicated in **FIG. 15A**. The mice were also treated with 3 mg/kg of the BMS 205820 polypeptide in PBS at the times indicated. Mice were sacrificed at day 18 and the presence of eosinophils assayed. As seen in **FIG. 15B**, the BMS 205820 polypeptide significantly inhibited the infiltration of eosinophils into the lungs, a measure of the asthmatic state.

Thus, inhibitors of nuclear translocation of cellular proteins have been disclosed. Although preferred embodiments of the subject invention have been described in some detail, it is understood that obvious variations can be made without departing from the spirit and the scope of the invention as defined by the appended claims.

10

15

20

25

30

35

WE CLAIM:

1. An isolated polypeptide comprising:
  - (1) a signal sequence peptide capable of  
5 delivering the polypeptide through the cytoplasmic membrane into a cell; and
  - (2) at least two nuclear localization sequences (NLSs),  
wherein said polypeptide is capable of  
10 inhibiting nuclear translocation of a cellular protein.
2. The polypeptide of claim 1, wherein the  
15 signal sequence is interchangeably flanked at its amino- and carboxy-termini by first and second NLSs.
3. The polypeptide of either of claims 1  
or 2, wherein the signal sequence peptide is the  
antennapedia homeodomain signal sequence peptide, the  
20 fibroblast growth factor signal sequence peptide, the human immunodeficiency virus (HIV) Tat signal sequence peptide, or the Hsc70 signal sequence peptide.
4. The polypeptide of claim 3, wherein the  
25 signal sequence peptide is the antennapedia homeodomain signal sequence peptide.
5. The polypeptide of claim 4, wherein the  
30 signal sequence peptide comprises the amino acid sequence RQIKIWFQNRRMKWKK (SEQ ID NO:7).
6. The polypeptide of claim 3, wherein the  
35 signal sequence peptide is the fibroblast growth factor signal sequence peptide.

7. The polypeptide of claim 6, wherein the signal sequence peptide comprises the amino acid sequence AAVALLPAVLLALLA (SEQ ID NO:8).

5 8. The polypeptide of claim 3, wherein the signal sequence peptide is the HIV Tat signal sequence peptide.

10 9. The polypeptide of claim 8, wherein the signal sequence peptide comprises the amino acid sequence CFITKALGISYGRKKRRQRRRPQGSQTH (SEQ ID NO:9).

15 10. The polypeptide of either of claims 1 or 2, wherein the first and second NLSSs may be the same or different and are peptides comprising the amino acid sequence selected from the group consisting of PKKKRKV (SEQ ID NO:10), KKKRKVC (SEQ ID NO:11), GKKRSKA (SEQ ID NO:12), KRPRP (SEQ ID NO:13), GNKAKRQRST (SEQ ID NO:14), GGAAKRVKLD (SEQ ID NO:15),  
20 SALIKKKKKMAP (SEQ ID NO:16), RKLKKLGN (SEQ ID NO:17), PPKKKP (SEQ ID NO:18), ASKSRKRKL (SEQ ID NO:19), KKKYK (SEQ ID NO:20), KKKYKC (SEQ ID NO:21), KSKKK (SEQ ID NO:22), KRVKLC (SEQ ID NO:23), and AKRVKL (SEQ ID NO:6).

25 11. The polypeptide of claim 7, wherein the first and second NLSSs may be the same or different and are peptides comprising the amino acid sequence selected from the group consisting of PKKKRKV (SEQ ID  
30 NO:10), KKKRKVC (SEQ ID NO:11), GKKRSKA (SEQ ID NO:12), KRPRP (SEQ ID NO:13), GNKAKRQRST (SEQ ID NO:14), GGAAKRVKLD (SEQ ID NO:15), SALIKKKKKMAP (SEQ ID NO:16), RKLKKLGN (SEQ ID NO:17), PPKKKP (SEQ ID NO:18), ASKSRKRKL (SEQ ID NO:19), KKKYK (SEQ ID  
35 NO:20), KKKYKC (SEQ ID NO:21), KSKKK (SEQ ID NO:22), KRVKLC (SEQ ID NO:23), and AKRVKL (SEQ ID NO:6).

12. The polypeptide of claim 11, wherein the first and second NLSs comprise the amino acid sequence PKKKRKV (SEQ ID NO:10).

5           13. The polypeptide of claim 11, wherein the first and second NLSs comprise the amino acid sequence PKKKRKV (SEQ ID NO:10) and KKKRKVC (SEQ ID NO:11), respectively, KKKYK (SEQ ID NO:20) and KKKYKC (SEQ ID NO:21), respectively, or AKRVKL (SEQ ID NO:6) and KRVKLC (SEQ ID NO:23), respectively.

14. The polypeptide of claim 12 comprising the amino acid sequence PKKKRKVAAVALLPAVLLALLAPKKKKRKV (SEQ ID NO:24).

15           15. The polypeptide of claim 13 comprising the amino acid sequence PKKKRKVAAVALLPAVLLALLAPKKKKRKVC (SEQ ID NO:1).

20           16. The polypeptide of any of claims 1-15, wherein the cellular protein is a transcription factor.

25           17. The polypeptide of claim 16, wherein the transcription factor is NF- $\kappa$ B.

18. A method of introducing an exogenous polypeptide comprising a nuclear localization sequence (NLS) into an intact cell so as to inhibit nuclear translocation of a cellular protein, comprising:

30           (a) providing a polypeptide according to any of claims 1-17; and

            (b) contacting the cell with the polypeptide for a period of time effective to  
35 introduce the exogenous polypeptide into the cell.

19. A method of suppressing an immune response in a subject comprising administering to the subject an immunosuppressive amount of a polypeptide according to any of claims 1-17.

5

20. A method of treating or preventing a viral infection in a subject, comprising administering to the individual an effective antiviral amount of a polypeptide according to any of claims 1-17.

10

21. A method of transcriptionally modulating the expression of cellular genes comprising, contacting a target cell with an inhibitor of nuclear translocation of a cellular protein, wherein the cellular protein is a transcription factor, and wherein the inhibitor comprises a polypeptide according to any of claims 1-17.

15

22. Use of a polypeptide according to any of claims 1-17 for the manufacture of a composition useful for suppressing an immune response in a subject.

20

23. Use of a polypeptide according to any of claims 1-17 for the manufacture of a composition useful for treating or preventing a viral infection in a subject.

25

24. Use of a polypeptide according to any of claims 1-17 for the manufacture of a composition useful for transcriptionally modulating the expression of cellular genes.

30

35



1/19




FIG. 1A

2/19




FIG. 1B

3/19




FIG. IC

4/19




FIG. 2

5/19




FIG. 3

6/19




FIG. 4A

7/19




FIG. 4B

8/19




FIG. 5A



9/19




FIG. 5B

10/19




FIG. 6A




FIG. 6B

11/19




FIG. 6C




FIG. 7

12/19

SV40MEM peptide

NF- $\kappa$ B MEM peptide




FIG. 8A




FIG. 8B

13/19




FIG. 9




FIG. 10

14/19




FIG. 11A

15/19





FIG. 11B




16/19



Treatment

FIG. 12



**FIG. 13A****FIG. 13B****FIG. 13C**

18/19




FIG. 14




FIG. 15A




FIG. 15B

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/16217

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :A61K 38/02, 38/04, 38/16; C07K 7/06, 7/08  
US CL :514/2; 530/300, 326, 327

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/2; 530/300, 326, 327

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
SEQ Search, APS, Dialog

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94/04696 A1 (MILES, INC.) 03 March 1994 (03-03-94), see especially the abstract and pages 7-8.	1-15
X, P	BERLOSE et al. Conformational and associative behaviours of the third helix of antennapedia homeodomain in membrane-mimetic environments. European Journal of Biochemistry. December 1996, Vol. 242, pages 372-386, especially the introduction and the discussion.	1-15

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* *A* *B* *L* *O* *P*	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	*T* *X* *Y* *A*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
--------------------------------------	---	--------------------------	--

Date of the actual completion of the international search

03 DECEMBER 1997

Date of mailing of the international search report

14 JAN 1998

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

WILLIAM SANDALS

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/16217

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KENTRUP et al. Dyrk, a dual specificity protein kinase with unique structural features whose activity is dependent on tyrosine residues between subdomains VII and VIII. The Journal of Biological Chemistry. 16 February 1996, Vol. 271, No. 7, pages 3488-3495, see especially the abstract.	1-15
Y	MUKAIGAWA et al. Two signals mediate nuclear localization of influenza virus (A/WSN/33) polymerase basic protein 2. Journal of Virology. January 1991, Vol. 65, No. 1, pages 254-253, see entire article.	1-15
Y	HINKES et al. Organization and promoter activity of the mouse syndecan-1 gene. Journal of Biological Chemistry. 25 May 1993, Vol. 268, No. 15, pages 11440-11448, see especially the abstract and introduction.	1-15
Y	IMAMURA et al. Identification of a heparin-binding growth factor-1 nuclear translocation sequence by deletion mutation analysis. Journal of Biological Chemistry. 15 March 1992, Vol. 267, No. 8, pages 5676-5679, see especially the abstract.	1-15
Y	PEREZ et al. Antennapedia homeobox as a signal for the cellular internalization and nuclear addressing of a small exogenous peptide. Journal of Cell Science. 17 August 1992, Vol. 102, pages 717-722, see the entire article.	1-15

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/16217

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 16-24  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐  
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**